

Nutrient dynamics in rhithron zone of Shivalik Himalayan stream Sahastradhara, Dehradun (Uttaranchal)

D. S. Malik and Pawan K. Bharti

Department of Zoology and Environmental science,
Gurukul Kangri University, Harwar-249404 (Uttaranchal).

Abstract

Sahastradhara hill-stream is a major tributary of river Song, 15 Km. far from Dehradun. It has a great importance as a picnic tourist spot of the region, whereas there is no more industrial and agricultural pollution. Hill streams of the area are the cradle of nutrients especially calcium and magnesium in the form of ions. Anthropogenic factors mainly tourists activities and catchments runoff may influence the index of nutrients in stream water. Nutrients play a vital role into niche restoration maintenance, self-regulation and water quality. Location variation in nutrients concentration of the stream was studied with special reference to correlation with physico-chemical parameters of the stream water. Calcium and Magnesium were observed in higher range in Sahastradhara stream in comparison to other hill-streams. Calcium and Magnesium were found maximum 104.21 mg/l and 68 mg/l respectively in summer season.

Key words: Nutrient dynamics, Sahastradhara stream, Rhithron zone, Water quality.

Introduction

The Himalayas are the cradle of a large number of streams and mighty rivers. Lesser Himalaya consists of a huge part of Uttaranchal. Dehradun, Harwar & Rishikesh are the main religious and beautiful cities, which are situated in the foothills of Garhwal Himalaya. A hill-stream Sahastradhara, 15 Km. far from Dehradun city is the study site for the accounting of nutrients in natural water and fluctuations at the different locations along with the stream. Adequate understanding of the Himalayan proglacial streams is extremely important for the development of a realistic program for utilizing the potential of water that exist in the form of snow and ice in the area.

The present study reveals to characteristics of the water nutrient chemistry, influenced by tourist's activity & quarrying of the geologically sedimental environments and to determine the nature and degree of anthropogenic impacts on qualitative & quantitative variations occurred in nutrients in relation to physico-chemical parameters of stream.

Materials and methods

The water samples were collected monthly from sampling site Sahastradhara stream from different sampling stations, Site-I, Site-II, Site-III, Site-IV, and Site-V during December, 2003 to May, 2004 in morning period 9:00 A. M. to 10:00 A. M.

The samples for nutrients and physico-chemical parameters were collected and analyzed by using rinsed borosil glassware, with the help of the procedure described by APHA (1995), Trivedi & Goel (1984), Santhanam *et al.* (1989).

Results & Discussion

Sahastradhara stream is flowing throughout a valley of Dehradun mountain chain in foothills of Shivalik Himalaya, enriched with limestone and lignite rocks, which affect the water quality of stream according to the locations. Nutrients concentration and related physico-chemical parameters from selected five sites are depicted in tables 1 & 2.

Sahastradhara stream has the spatio-temporal variations of water temperature, which plays a vital role in all physico-biochemical reactions and self-purification power of aquatic system (Badola & Singh, 1981). Higher value of temperature was found 18 °C in summer and minimum 12 °C in winter season. A direct relationship was established between the water temperature and free carbon dioxide (Hynes, 1970). Turbidity is striking characteristic of the physical status of the water bodies. Although in Sahastradhara stream water is clear because there is no more pollution, siltation was the main source of turbidity in tributaries. Detritus and other non-organic material being added to water mass due to rainfall and tourist activities (Camron, 1996). Maximum turbidity was recorded 7.0 JTU during rainy days and minimum 1.0 JTU in December. The maximum depth of photic zone provides the better biological production for all aquatic organisms.

Total solids were found in the range of 955 mg/l to 1535 mg/l towards summer season, due to the gradual increases in velocity of river which favoured effective sedimentation (Subramanian, 1979). Chemical oxygen demand represents chemically oxidizable organic matter load in water, while biochemical oxygen demand is only biodegradable materials. In the present study high values observed during summer months may be attributed maximum biological activities and high temperature, stimulate the growth of microorganisms (William *et al.*, 1993).

The pH of natural water was controlled in a great extent by the interaction of hydroxyl ions arising from the hydrolysis of bicarbonate (Sharma, 1986). The pH of Sahastradhara stream was recorded slightly alkaline (7.2 - 8.1). Hardness is mainly due to percentage of calcium and magnesium salts of bicarbonates, carbonates, sulphates and chlorides, while the value of alkalinity occurred due to presence of bicarbonates. The highest concentration of hardness was analyzed 250 mg/l during warm months. A positive relationship between hardness and alkalinity was recorded in river Ganga at Rishikesh (Chopra and Patrick, 1994).

Maximum chloride concentration was recorded (18.46 mg/l) in the month of May and minimum in December (11.36 mg/l). Chloride and hardness showed a positive relationship to one another (Chopra and Patric, 1994). Chloride was found in the form of chloride ion, and one of the major inorganic anion present in natural water.

Calcium and magnesium the dominant cations, and these represent the main weathering products, but significant hydro-chemical differences between the five sampling sites associated with the bedrock geology exist (Jenkins *et al.*, 1995). Calcium is one of the essential nutrients, which plays an important role in biological system. Maximum calcium concentration was recorded (104.208 mg/l) in the month of May and minimum in January (77.154 mg/l). Calcium concentrations were found maximum about 7.5% of the total solids present in the stream water. Positive relationship between, calcium and temperature was also reported by Khanna and Singh (2000) in river Suswa, Dehradun. Magnesium is also an essential element but it is toxic at higher concentration. The amount of magnesium in Sahastradhara was found maximum (68.0 mg/l) in the month of April and minimum (45.0 mg/l) in December.

During the summer season nutrients concentration in hill-streams became more. Miller *et al.* (1997) described the nutrients availability in selected environmental settings of the Potomac River and Cameron, E. M. (1996) showed the similar type of fluctuation in Fraser river of British Columbia. Sodium is one of the most common cations has no adverse effect on human health at lower concentration. Bond, H.B. (1979) described similar nutrients concentration pattern in a stream draining a mountain ecosystem in Utah. Maximum sodium concentration in Sahastradhara was recorded (11.0 mg/l) in the month of May and minimum in January (6.0 mg/l).

Potassium is naturally occurring element, released by the clay minerals, weathering and leaching from growing vegetation and decomposition of organic matter (Berndtsson, 1990). Maximum potassium concentration was recorded (18.0 mg/l) in the month of April and minimum in December (14.0 mg/l). Pande and Mishra (2000) also observed similar results on Sahastradhara hill-stream Dehradun. Relationship between nutrients and other physico-chemical parameters are given in the table-3 and percentages of nutrients among the total solids are given in table-4, during the study period. These high concentrations confirm the location of the calcareous formation associated with the Central Himalaya thrust. Similarly, samples from stream draining the slope of Doon valley, exhibit high concentrations of Mg relative to K and Na. Again this can presumably be attributed to local geological features. In general, the stream at site III with high Ca concentrations also exhibit relatively low K concentrations compared to the other sites.

The present results conclude that significant differences in stream water nutrient concentrations exist among different environmental settings within the five subunits. The environmental setting with the highest potential by more soluble nutrients, fluctuations in nutrient concentrations were the land use and carbonate bedrock that was predominated in the Doon valley especially, at main tourist spot & sulphur springs site III. Ca and Mg are the dominant cations in this alkaline tributary of river Song. Rock weathering is the most important mechanism controlling the water chemistry. The high concentrations of alkali earth metals, high alkalinity and the high (Ca+Mg)/(Na+K) ratio indicate that the water chemistry of the stream is controlled largely by carbonate weathering and partially by run-off process & tourists activities. The spatial variations in TDS are attributed to climatic and lithological control over the ionic concentrations.

References

- APHA (1995): Standard methods for examination of water and waste water. *American Public Health Association, 19th edition. Inc, New York*. pp:1150.
- Alan Jenkins; Jeremy Wilkinson and Robert Collins (1995): Hydro-chemical characteristics of headwater stream in the middle hills and high mountains of the Himalayas, Nepal. *Atmospheric Hydrologic and Ecological interaction- Part C. International conference on ecohydrology of high mountain areas. pp: 255-263.*
- Badola, S.P. and Singh, H.R. (1981) Hydrobiology of the river Alaknanda of Garhwal Himalaya. *Indian J. Ecol.*, 8(2): 269-276.
- Berndtsson, R. (1990): Transport and sedimentation of pollutants in a river reach: A chemical mass balance approach. *Wat. Resour. Res.* 26(7): 1549-1558.
- Bond, H. B. (1979): Nutrient concentrations patterns in a stream draining a montane ecosystem in Utah. *Ecology*. 60(6): 1184-1196.

Cameron, E. M. (1996): Hydrogeo-chemistry of the Fraser River British Columbia: Seasonal variation in major and minor components. *J. Hydrol.* 182(1-4): 209-255.

Chopra, A. K. and Patrick, N. J. (1994): Effect of domestic sewage on self-purification of Ganga water at Rishikesh. *A. Bio. Science.* 13(11): 75-82.

Hussain, S. I.; Subramanian, V. and Dhanpal, K. (1989): Chemical characteristics of meltwaters from a Himalaya glacier, India. *J. Hydrol.* 106: 98-106.

Hynes, H.B.N. (1970). The ecology of running waters. *Liverpool University Press, Liverpool, 4th impression:* 1-555.

Jenkins, A., Sloan W.T. & Cosby, B.J. (1995) Stream chemistry in the middle hills and high mountain of the Himalaya, Nepal. *Journal of Hydrology.* Vol-166, No-1-4: 61-79.

Khanna, D.R and Singh. R.K. (2000). Seasonal fluctuations in the plankton of Suswa River at Raiwala Dehradun. *Env. Conservation J.* 1(2 & 3): 89-92.

Miller, C. V.; Denis, J. M.; Ator, S. W. and Brakebill, J. W. (1997): Nutrients in stream during baseflow in selected environmental settings of the Potomac river basin. *J. American Wat. Resources Association.* 33(6): 1155-1171.

Odum, E. P. (1971): Fundamentals of Ecology. New York. W. B. Saunders and Co. pp:1-344.

Pande, R. K. and Asha Mishra (2000): Water quality study of freshwaters of Dehradun (Sahastradhara stream and Mussoorie Lake). *Aquacult.* 1: 57-62.

Santhanam, R.; Velaynatham, P. and Jegatheesan, G. (1989): A manual for fresh water ecology. *Daya Publishing House, Delhi- 110006:*1-92.

Shrama, R.C. (1986). Effect of physico-chemical factors on benthic fauna of Bhagirathi River Garhwal Himalaya. *Indian. J.Ecol.* 13(1): 133-137.

Subramanian, V. (1979) Chemical and suspended sediment characteristics of river of India. *J. Hydrol.* 44:37-55.

Trivedi, R. K. and Goel, P. K. (1984): Chemical and biological methods for water pollution studies. Karad. Environmental publication. pp: 1-25.

William, M.W., Brown, A. & Melack, J.M. (1993). Geochemical and hydrologic controls on the composition of surface water in the high elevation basin, *Sierra Nevada. Limnol. Oceanogr.* 38: 775-797.

Table- 1: Nutrients parameters (mean values in mg/l) of Sahastradhara stream.

Parameters	Site-I	Site-II	Site-III	Site-IV	Site-V
Chloride Mean	(11.36 - 15.62) 12.78±1.73	(11.36 - 16.58) 13.73±1.93	(12.78 - 18.46) 15.26±2.1	(9.94 - 18.46) 13.49±3.25	(8.52 - 16.58) 11.6±3.06
Calcium Mean	(80.16-88.17) 84.67±3.58	(80.16-92.18) 85.67±4.28	(77.15-104.20) 91.68±10.13	(79.15-99.19) 89.18±8.26	(80.16-96.19) 88.43±6.19
Magnesium Mean	(45 - 53) 48.66±3.09	(55 - 62) 57.33±3.29	(63.0 - 68.0) 64.66±2.36	(52.0 - 28.0) 55.33±2.49	(45.0 - 55.0) 51.0±5.29
Sodium	(6 - 8.5) 7.375±0.96	(7.0 - 8.2) 7.725±0.506	(8.0 - 9.5) 8.75±0.75	(7.0 - 10.0) 8.87±1.14	(7.5 - 11.0) 9.25±1.35
Potassium	(14 - 16) 15.13±0.74	(14.5 - 17.0) 15.75±0.90	(16.0 - 17.5) 16.62±0.65	(15.5 - 17.5) 16.5±0.79	(16.0 - 18.0) 17.0±1.35

Nutrient dynamics in rhithron zone

Table- 2: Physico-chemical parameters (mean values) of Sahastradhara stream.

Parameters	Site- I	Site- II	Site- III	Site- IV	Site- V
Temperature (°C)	(14 - 18)	(13 - 17.5)	(13 - 17)	(13.5 - 18)	(12 - 18)
Mean	15.75±1.14	15.37±1.63	15.0±1.41	15.75±1.6	15.75±2.277
Velocity (m/s)	(0.3 - 0.4)	(0.28 - 0.32)	(0.8 - 1.0)	(0.78 - 0.82)	(0.6 - 0.7)
Mean	0.3±0.14	0.9±0.1	0.35±0.05	0.8±0.02	0.65±0.05
Turbidity (JTU)	(1.0 - 5.0)	(2.0 - 5.0)	(3.0- 6.0)	(2.0 - 6.0)	(3.0- 7.0)
Mean	2.75±1.47	3.25±1.30	4.5±1.15	3.75±1.48	4.75±1.48
Total solids (mg/l)	(955 - 1427)	(1090 - 1332)	(1115 - 1443)	(1135 - 1520)	(1148 - 1535)
Mean	1218±183.2	1234.2±89.8	1277.2±122.6	1305.7±166.1	1290.7±146.1
TDS (mg/l)	(840 - 1278)	(895 - 1217)	(1010 - 1322)	(1039 - 1380)	(965 - 1420)
Mean	1093.25±174.9	1094.25±120.5	1140.5±134.6	1189±158.01	1134.25±176.1
TSS (mg/l)	(113 - 149)	(119 - 195)	(105 - 207)	(105 - 140)	(115 - 211)
Mean	124.75±14.39	140.0±28.16	136.75±40.95	116.75±13.8	156.5±41.69
pH	(7.2 - 8.1)	(7.3 - 8.1)	(7.5 - 7.9)	(7.6 - 8.0)	(7.2 - 8.1)
Mean	7.625±0.43	7.6±0.28	7.75±0.15	7.75±1.66	7.6±0.339
Free CO ₂ (mg/l)	(1.40 - 1.69)	(1.62 - 1.92)	(1.62 - 2.20)	(1.65 - 1.85)	(1.62 - 2.20)
Mean	1.6525±0.39	1.75±.017	1.825±0.25	1.74±0.09	1.87±0.21
D O (mg/l)	(7.04 - 11.08)	(6.84 - 10.07)	(7.04 - 9.47)	(6.04 - 9.06)	(6.04 - 9.47)
Mean	8.41±1.62	8.46±1.18	8.65±0.95	7.55±0.87	7.49±1.36
BOD (mg/l)	(1.60 - 2.0)	(1.7 - 2.2)	(2.0 - 2.5)	(1.8 - 2.2)	(1.9 - 2.3)
Mean	1.875±0.162	1.98±0.23	2.25±.18	2.0±0.14	2.05±0.15
COD (mg/l)	(2.1 - 3.0)	(2.2 - 2.9)	(2.6 - 3.2)	(2.4 - 3.0)	(2.4 - 2.9)
Mean	2.55±0.32	2.6±0.25	2.9±0.22	2.68±0.22	2.7±0.18
Hardness (mg/l)	(190 - 210)	(185 - 220)	(200 - 250)	(180 - 230)	(185 - 235)
Mean	196.25±8.2	200.0±15.41	222.5±19.2	200.0±18.71	198.75±21.02
Alkalinity (mg/l)	(150 - 480)	(140 - 500)	(170 - 600)	(130 - 570)	(120 - 540)
Mean	250.0±134.7	262.5±140.8	305±172.1	292.5±168.8	317.5±175.2

Table- 3: Correlation (=CORREL) between Nutrients (mean values) and Physicochemical parameters (mean values) of Sahastradhara stream.

Parameters	Chloride	Calcium	Magnesium	Sodium	Potassium
Temperature (°C)	-0.86097	-0.51833	-0.92056	-0.01186	-0.14275
Velocity (m/s)	-0.17595	-0.12077	0.007842	0.179873	0.225839
Turbidity (JTU)	0.012697	0.818834	0.400371	0.92284	0.959357
Total solids (mg/l)	-0.00448	0.788427	0.273292	0.949891	0.906666
T. D. S. (mg/l)	0.133154	0.71132	0.25802	0.764443	0.682478
T. S. S. (mg/l)	-0.35586	0.096981	0.004271	0.35689	0.46296
pH (pH unit)	0.681329	0.761113	0.623067	0.395679	0.345766
Free CO ₂ (mg/l)	-0.03224	0.686443	0.396104	0.833271	0.90927
D.O. (mg/l)	0.679888	-0.12231	0.471533	-0.65112	-0.55633
B. O. D. (mg/l)	0.607899	0.912138	0.845587	0.615583	0.694262
C. O. D. (mg/l)	0.595229	0.954861	0.801927	0.654354	0.704159
Hardness (mg/l)	0.814816	0.795954	0.897315	0.312648	0.386789
Alkalinity (mg/l)	-0.04111	0.829565	0.331085	0.97707	0.985243

Table- 4: Percentage of different nutrients among the total solids at different location of Sahastradhara stream at Dehradun.

Nutrients	Site-I	Site-II	Site-III	Site-IV	Site-V
Total solids	100%	100%	100%	100%	100%
Chloride	1.0492%	1.1125%	1.1948%	1.0331%	0.8987%
Calcium	6.951%	6.941%	7.178%	6.830%	6.851%
Magnesium	3.995%	4.645%	5.062%	4.238%	3.951%
Sodium	0.6055%	0.6263%	0.6852%	0.6793%	0.7166%
Potassium	1.242%	1.276%	1.301%	1.264%	1.317%